NICE National Institute for Health and Care Excellence

Coronavirus (COVID-19)

For information on how NICE is supporting the NHS and social care, view our new rapid guidelines and evidence reviews

(https://web.archive.org/web/20200331052542/https://www.nice.org.uk/coronavirus).

Learn about the government response to coronavirus on GOV.UK

(https://web.archive.org/web/20200331052542/https://gov.uk/coronavirus).

Guidance (/web/20200331092821/https://www.nice.org.uk/guidance/ng159)

Tools and resources

(/web/20200331092821/https://www.nice.org.uk/guidance/ng159/resources)

Overview (/web/20200331092821/https://www.nice.org.uk/guidance/ng159)

1 Admission to hospital

2 Admission to critical care

(/web/20200331092821/https://www.nice.org.uk/guidance/ng159/chapter/2-Admission-to-critical-care)

3 Starting, reviewing and stopping critical care treatment

(/web/20200331092821/https://www.nice.org.uk/guidance/ng159/chapter/3-Starting-reviewing-and-stopping-critical-care-treatment)

4 Clinical decision-making

(/web/20200331092821/https://www.nice.org.uk/guidance/ng159/chapter/4-Clinical-decision-making)

5 Service organisation

(/web/20200331092821/https://www.nice.org.uk/guidance/ng159/chapter/5-Service-organisation)

Update information

(/web/20200331092821/https://www.nice.org.uk/guidance/ng159/chapter/Update-information)

Guidance

1 Admission to hospital

- 1.1 On admission to hospital, assess all adults for frailty, irrespective of COVID-19 status.
 - Use the <u>Clinical Frailty Scale</u>
 (https://web.archive.org/web/20200331092821/https://static1.squarespace.com/static (CFS), available from the NHS Specialised Clinical Frailty Network
 (https://web.archive.org/web/20200331092821/https://www.scfn.org.uk/), as part of a holistic assessment.
 - Be aware of the limitations of using the CFS as the sole assessment of frailty. The
 CFS should not be used in younger people, people with stable long-term
 disabilities (for example, cerebral palsy), learning disabilities or autism. An
 individualised assessment is recommended in all cases where the CFS is not
 appropriate.
 - Consider comorbidities and underlying health conditions in all cases.
 - Record the frailty assessment in the patient's medical record.
 [amended 25 March 2020]
- 1.2 When patients with possible COVID-19 have been identified, follow appropriate <u>UK</u> government guidance on infection prevention and control measures (https://www.gov.uk/government/publicat novel-coronavirus-infection-prevention-and-control). This includes recommendations on:
 - patient transfers and transport
 - segregation and cohorting
 - personal protective equipment
 - aerosol-generating procedures.
- 1.3 If COVID-19 is diagnosed in someone not isolated from admission or presentation, follow UK government guidance on actions required when a case was not diagnosed on admission

(https://web.archive.org/web/20200331092821/https://www.gov.uk/government/publicat 19-guidance-for-healthcare-providers-who-have-diagnosed-a-case-within-their-facility/covid-19-actions-required-when-a-case-was-not-diagnosed-on-admission).